


# CLAIMS

The subject matter claimed is:

1. A composition of matter having a formula represented by



5 wherein x is an integer of about 8 to about 1,200, X is a linker, and Y is a residue of a sterol comprising a 3-ol group.

2. The composition of matter of claim 1 wherein x is about 581.


3. The composition of matter of claim 1 wherein X is  
10 -O-CO-.

4. The composition of matter of claim 1 wherein Y is a cholesterol residue.

5. The composition of matter of claim 1 wherein Y is a member selected from the group consisting of residues of  
15 cholesterol, cholestanol, coprosterol, epicholestanol, epicholesterol, ergostanol,  $\alpha$ -ergosterol,  $\beta$ -ergosterol,  $\gamma$ -ergosterol, ergosterol, 22,23-dihydroergosterol, stigmaterol, stigmastanol, (3 $\beta$ )-7-dehydrocholesterol, desmosterol, allocholesterol, 24-hydroxycholesterol, 25-hydroxycholesterol,  
20 campesterol,  $\alpha_1$ -sitosterol,  $\beta$ -sitosterol,  $\gamma$ -sitosterol,

lumisterol, pyrocalciferol, isopyrocalciferol, azacosterol,  
neoergosterol, and dehydroergosterol.


6. A composition of matter having a formula represented by



5 wherein x is an integer of about 8 to about 1,200, and Y is a  
cholesterol residue.

7. The composition of matter of claim 6 wherein x is about  
581.

8. A composition of matter having a formula represented by



15 wherein x is an integer of about 0 to about 1,200, y is an  
integer of about 0 to about 1,200, with the proviso that x + y is  
about 8 to about 1,200, X is a linker, and Y is a residue of a  
sterol comprising a 3-ol group.


9. The composition of matter of claim 8 wherein x + y is  
20 about 581.

10. The composition of matter of claim 8 wherein X is -CO-.

11. The composition of matter of claim 8 wherein Y is a cholesterol residue.

12. The composition of matter of claim 8 wherein Y is a member selected from the group consisting of residues of  
5 cholesterol, cholestanol, coprosterol, epicholestanol, epicholesterol, ergostanol,  $\alpha$ -ergosterol,  $\beta$ -ergosterol,  $\gamma$ -ergosterol, ergosterol, 22,23-dihydroergosterol, stigmasterol, stigmasterol, (3 $\beta$ )-7-dehydrocholesterol, desmosterol, allocholesterol, 24-hydroxycholesterol, 25-hydroxycholesterol,  
10 campesterol,  $\alpha_1$ -sitosterol,  $\beta$ -sitosterol,  $\gamma$ -sitosterol, lumisterol, pyrocalciferol, isopyrocalciferol, azacosterol, neoergosterol, and dehydroergosterol.


13. A composition of matter having a formula represented by



wherein x is an integer of about 0 to about 1,200, y is an  
20 integer of about 0 to about 1,200, with the proviso that x + y is about 8 to about 1,200, and Y is a cholesterol residue.

14. The composition of matter of claim 13 wherein x + y is about 581.

15. A composition of matter having a formula represented by



wherein x is an integer of about 0 to about 1,200, y is an integer of about 0 to about 1,200, with the proviso that x + y is about 8 to about 1,200, X<sub>1</sub> and X<sub>2</sub> are linkers, and Y<sub>1</sub> and Y<sub>2</sub> are residues of a sterol comprising a 3-ol group.

16. The composition of matter of claim 15 wherein x + y is about 581.


17. The composition of matter of claim 15 wherein X<sub>1</sub> is -O-CO- and X<sub>2</sub> is -CO-.

18. The composition of matter of claim 15 wherein Y<sub>1</sub> and Y<sub>2</sub> are cholesterol residues.

19. The composition of matter of claim 15 wherein Y<sub>1</sub> and Y<sub>2</sub> are members independently selected from the group consisting of residues of cholesterol, cholestanol, coprosterol, epicholestanol, epicholesterol, ergostanol, α-ergosterol, β-ergosterol, γ-ergosterol, ergosterol, 22,23-dihydroergosterol, stigmasterol, stigmastanol, (3β)-7-dehydrocholesterol,


desmosterol, allocholesterol, 24-hydroxycholesterol, 25-hydroxycholesterol, campesterol,  $\alpha_1$ -sitosterol,  $\beta$ -sitosterol,  $\gamma$ -sitosterol, lumisterol, pyrocalciferol, isopyrocalciferol, azacosterol, neoergosterol, and dehydroergosterol.

20. A composition of matter having a formula represented by



wherein x is an integer of about 0 to about 1,200, y is an integer of about 0 to about 1,200, with the proviso that x + y is about 8 to about 1,200, and Y is a cholesterol residue.

21. A complex comprising a mixture of a nucleic acid and a composition of matter having a formula represented by



wherein x is an integer of about 8 to about 1,200, X is a linker, and Y is a residue of a sterol comprising a 3-ol group.

22. The complex of claim 21 wherein x is about 581.

23. The complex of claim 21 wherein X is -O-CO-.

24. The complex of claim 21 wherein Y is a cholesterol


residue.

25. The complex of claim 21 wherein X is -O-CO- and Y is a cholesterol residue.

26. The complex of claim 21 wherein Y is a member selected  
5 from the group consisting of residues of cholesterol,  
cholestanol, coprosterol, epicholestanol, epicholesterol,  
ergostanol,  $\alpha$ -ergosterol,  $\beta$ -ergosterol,  $\gamma$ -ergosterol,  
ergosterol, 22,23-dihydroergosterol, stigmasterol, stigmastanol,  
10 (3 $\beta$ )-7-dehydrocholesterol, desmosterol, allocholesterol, 24-  
hydroxycholesterol, 25-hydroxycholesterol, campesterol,  $\alpha_1$ -  
sitosterol,  $\beta$ -sitosterol,  $\gamma$ -sitosterol, lumisterol,  
pyrocalciferol, isopyrocalciferol, azacosterol, neoergosterol,  
and dehydroergosterol.

27. The complex of claim 21 wherein the nucleic acid  
15 comprises a plasmid.

28. A complex comprising a mixture of a nucleic acid and a  
composition of matter having a formula represented by:



wherein x is an integer of about 0 to about 1,200, y is an integer of about 0 to about 1,200, with the proviso that x + y is about 8 to about 1,200, X is a linker, and Y is a residue of a sterol comprising a 3-ol group.

5           29. The complex of claim 28 wherein x + y is about 581.

30. The complex of claim 28 wherein X is -CO-.

31. The complex of claim 28 wherein Y is a cholesterol residue.


10           32. The complex of claim 28 wherein X is -CO- and Y is a cholesterol residue.

33. The complex of claim 28 wherein Y is a member selected from the group consisting of residues of cholesterol, cholestanol, coprosterol, epicholestanol, epicholesterol, ergostanol,  $\alpha$ -ergosterol,  $\beta$ -ergosterol,  $\gamma$ -ergosterol, 15 ergosterol, 22,23-dihydroergosterol, stigmasterol, stigmastanol, (3 $\beta$ )-7-dehydrocholesterol, desmosterol, allocholesterol, 24-hydroxycholesterol, 25-hydroxycholesterol, campesterol,  $\alpha_1$ -sitosterol,  $\beta$ -sitosterol,  $\gamma$ -sitosterol, lumisterol, pyrocalciferol, isopyrocalciferol, azacosterol, neoergosterol,

and dehydroergosterol.

34. The complex of claim 28 wherein the nucleic acid comprises a plasmid.

34. A complex comprising a mixture of a nucleic acid and a composition of matter having a formula represented by



wherein x is an integer of about 0 to about 1,200, y is an integer of about 0 to about 1,200, with the proviso that x + y is about 8 to about 1,200, X<sub>1</sub> and X<sub>2</sub> are linkers, and Y<sub>1</sub> and Y<sub>2</sub> are residues of a sterol comprising a 3-ol group.

35. The complex of claim 34 wherein x + y is about 581.

36. The complex of claim 34 wherein X<sub>1</sub> is -O-CO- and X<sub>2</sub> is -CO-.

37. The complex of claim 34 wherein Y<sub>1</sub> and Y<sub>2</sub> are cholesterol residues.



38. The complex of claim 34 wherein  $X_1$  is  $-O-CO-$ ,  $X_2$  is  $-CO-$ , and  $Y_1$  and  $Y_2$  are cholesterol residues.

39. The complex of claim 34 wherein  $Y_1$  and  $Y_2$  are members independently selected from the group consisting of residues of  
5 cholesterol, cholestanol, coprosterol, epicholestanol, epicholesterol, ergostanol,  $\alpha$ -ergosterol,  $\beta$ -ergosterol,  $\gamma$ -ergosterol, ergosterol, 22,23-dihydroergosterol, stigmasterol, stigmasterol, (3 $\beta$ )-7-dehydrocholesterol, desmosterol, allocholesterol, 24-hydroxycholesterol, 25-hydroxycholesterol,  
10 campesterol,  $\alpha_1$ -sitosterol,  $\beta$ -sitosterol,  $\gamma$ -sitosterol, lumisterol, pyrocalciferol, isopyrocalciferol, azacosterol, neoergosterol, and dehydroergosterol.

40. The complex of claim 34 wherein the nucleic acid comprises a plasmid.

15 41. A method of making an L-shaped linear polyethylenimine sterol conjugate comprising reacting a linear polyethylenimine having an average molecular weight of about 423 to about 50,000 and comprising a terminal hydroxyl group with a chloroformate ester of a sterol comprising a 3-ol group, thereby resulting in  
20 the L-shaped linear polyethylenimine sterol conjugate comprising the sterol covalently bonded to the terminal hydroxyl group.

42. The method of claim 41 wherein the sterol is a member selected from the group consisting of cholesterol, cholestanol, coprosterol, epicholestanol, epicholesterol, ergostanol,  $\alpha$ -ergostenol,  $\beta$ -ergostenol,  $\gamma$ -ergostenol, ergosterol, 22,23-  
5 dihydroergosterol, stigmasterol, stigmastanol, (3 $\beta$ )-7-dehydrocholesterol, desmosterol, allocholesterol, 24-hydroxycholesterol, 25-hydroxycholesterol, campesterol,  $\alpha_1$ -sitosterol,  $\beta$ -sitosterol,  $\gamma$ -sitosterol, lumisterol, pyrocalciferol, isopyrocalciferol, azacosterol, neoergosterol,  
10 and dehydroergosterol.

43. The method of claim 41 wherein the sterol comprises cholesterol.

44. A method of making a T-shaped linear polyethylenimine sterol conjugate comprising:

15 (a) reacting a linear polyethylenimine having an average molecular weight of about 423 to about 50,000 and comprising a terminal hydroxyl group and a plurality of secondary amine nitrogen atoms with a protecting reagent such that the protecting reagent bonds with the terminal hydroxyl group, resulting in a  
20 protected linear polyethylenimine;

(b) reacting a chloroformate ester of a sterol comprising a 3-ol group with the protected linear polyethylenimine such that

the chloroformate ester of a sterol bonds with at least one of the plurality of secondary amine nitrogen atoms, resulting in a protected T-shaped linear polyethylenimine sterol conjugate; and

(c) deprotecting the protected T-shaped linear  
5 polyethylenimine sterol conjugate with a deprotecting reagent, resulting in the T-shaped linear polyethylenimine sterol conjugate.

45. The method of claim 44 wherein the sterol is a member selected from the group consisting of cholesterol, cholestanol,  
10 coprosterol, epicholestanol, epicholesterol, ergostanol,  $\alpha$ -ergostenol,  $\beta$ -ergostenol,  $\gamma$ -ergostenol, ergosterol, 22,23-dihydroergosterol, stigmasterol, stigmastanol, (3 $\beta$ )-7-dehydrocholesterol, desmosterol, allocholesterol, 24-hydroxycholesterol, 25-hydroxycholesterol, campesterol,  $\alpha_1$ -  
15 sitosterol,  $\beta$ -sitosterol,  $\gamma$ -sitosterol, lumisterol, pyrocalciferol, isopyrocalciferol, azacosterol, neoergosterol, and dehydroergosterol.

46. The method of claim 44 wherein the sterol comprises cholesterol.

20 47. The method of claim 44 wherein the protecting reagent comprises chlorotrimethylsilane and the protected linear

polyethylenimine comprises linear polyethylenimine trimethyl silane.

48. The method of claim 47 wherein the deprotecting reagent comprises trifluoroacetic acid.

5           49. A method of making an LT-shaped linear polyethylenimine sterol conjugate comprising reacting a linear polyethylenimine having an average molecular weight of about 423 to about 50,000 and comprising a terminal hydroxyl group and a plurality of secondary amine nitrogen atoms with a chloroformate ester of a sterol comprising a 3-ol group such that chloroformate ester of a sterol bonds with the terminal hydroxyl group and at least one of the plurality of secondary amine nitrogen atoms, thereby  
10           resulting in the LT-shaped linear polyethylenimine sterol conjugate.

15           50. The method of claim 49 wherein the sterol is a member selected from the group consisting of cholesterol, cholestanol, coprosterol, epicholestanol, epicholesterol, ergostanol,  $\alpha$ -ergostenol,  $\beta$ -ergostenol,  $\gamma$ -ergostenol, ergosterol, 22,23-dihydroergosterol, stigmasterol, stigmastanol, (3 $\beta$ )-7-  
20           dehydrocholesterol, desmosterol, allocholesterol, 24-hydroxycholesterol, 25-hydroxycholesterol, campesterol,  $\alpha_1$ -

sitosterol,  $\beta$ -sitosterol,  $\gamma$ -sitosterol, lumisterol,  
pyrocalciferol, isopyrocalciferol, azacosterol, neoergosterol,  
and dehydroergosterol.

51. The method of claim 49 wherein the sterol comprises  
5 cholesterol.

52. A method of delivering a nucleic acid into a mammalian  
cell comprising:

(a) mixing the nucleic acid with an L-shaped, T-shaped, or  
LT-shaped linear polyethylenimine sterol conjugate to result in a  
10 complex;

(b) contacting the mammalian cell with the complex such  
that the complex enters the mammalian cell, thereby delivering  
the nucleic acid into the mammalian cell.